

The Pascal Triangle of a Discrete Image: Definition, Properties and Application to Shape Analysis

Mireille Boutin*, Shanshan Huang†

Abstract

We define the Pascal Triangle of a discrete (gray scale) image as a pyramidal arrangement of complex-valued moments and we explore its geometric significance. In particular, we show that the entries of row k of this triangle correspond to the Fourier series coefficients of the moment of order k of the Radon transform of the image. Group actions on the plane can be naturally prolonged onto the entries of the Pascal Triangle. We study the prolongation of some common group actions, such as rotations and reflections, and we propose simple tests for detecting equivalences and self-equivalences under these group actions. The motivating application of this work is the problem of recognizing “shapes” on images, for example characters, digits or simple graphics.

1 Definition and Reconstruction Properties

Let $\{(x_k, y_k)\}_{k=1}^N$ with $x_k, y_k \in \mathbb{R}$ represent the pixel locations of a digital image. For simplicity, we use complex coordinates $z_k = x_k + iy_k$. Consider a gray scale image defined on $\{z_k\}_{k=1}^N$. More specifically, we have a mapping $\rho : \{z_k\}_{k=1}^N \rightarrow \mathbb{R}_{\geq 0}$, where $\rho(z_k)$ represents the intensity of pixel z_k ¹. Denote the discrete gray scale image by $I = \{(z_k, \rho(z_k))\}_{k=1}^N$.

Consider the following moment matrix:

$$\tau_N(I) = \begin{pmatrix} \mu_{0,0} & \mu_{1,0} & \mu_{2,0} & \mu_{0,3} & \cdots & \mu_{N-1,0} \\ \mu_{0,1} & \mu_{1,1} & \mu_{2,1} & \cdots & \cdots & \mu_{N-1,1} \\ \mu_{0,2} & \mu_{1,2} & \cdots & \cdots & \cdots & \mu_{N-1,2} \\ \mu_{0,3} & \cdots & \cdots & \cdots & \cdots & \mu_{N-1,3} \\ \vdots & & & & & \vdots \\ \mu_{0,N-1} & \mu_{1,N-1} & \cdots & \cdots & \cdots & \mu_{N-1,N-1} \end{pmatrix}_{N \times N},$$

*School of Electrical and Computer Engineering, Purdue University. Email: mboutin@purdue.edu

†Department of Mathematics, Purdue University. Email: huang94@purdue.edu

¹We use $\mathbb{R}_{\geq 0}$ instead of a specific discrete domain such as $\{0, 1, \dots, 255\}$ for more generality.

where

$$\mu_{j,l} = \sum_{k=1}^N z_k^j \bar{z}_k^l \rho(z_k), \quad j, l \in \mathbb{Z}_{\geq 0}$$

is the complex moment of order (j, l) for the discrete image I . Observe the conjugate symmetry property of the moments $\mu_{j,l} = \bar{\mu}_{l,j}$. In particular, $\mu_{j,j} \in \mathbb{R}$, $\forall j \in \mathbb{Z}_{\geq 0}$.

We can express the relationship between the moments and the image I in matrix form:

$$\tau_N(I) = Z^\dagger W Z, \quad (1)$$

where

$$Z = \begin{pmatrix} 1 & z_1 & z_1^2 & \cdots & z_1^{N-1} \\ 1 & z_2 & z_2^2 & \cdots & z_2^{N-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_N & z_N^2 & \cdots & z_N^{N-1} \end{pmatrix}, \quad W = \begin{pmatrix} \rho(z_1) & 0 & \cdots & 0 \\ 0 & \rho(z_2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \rho(z_N) \end{pmatrix},$$

and $Z^\dagger$ is the conjugate transpose of Z . Observe that Z is a Vandermonde matrix and therefore is invertible when the pixel locations z_k are pairwise distinct. Therefore, if the pixel coordinates are known and pairwise distinct, one can reconstruct the image I by matrix inversion: $W = (Z^{-1})^\dagger \tau(I) Z^{-1}$.

Definition 1. Let r be a nonnegative integer and let I be a discrete gray scale image. The Pascal Triangle $T^r(I)$ of order r of I is the following pyramid:

$$\begin{array}{ccccccc} & & & & \mu_{0,0} & & \\ & & & & & & \\ & & & & \mu_{0,1} & \mu_{1,0} & \\ & & & & \mu_{0,2} & 2\mu_{1,1} & \mu_{2,0} \\ & & & & \mu_{0,3} & 3\mu_{1,2} & 3\mu_{2,1} & \mu_{3,0} \\ & & & & \mu_{0,4} & 4\mu_{1,3} & 6\mu_{2,2} & 4\mu_{3,1} & \mu_{4,0} \\ & & & & & & \vdots & & \\ & & & & \mu_{0,r} & \binom{r}{1} \mu_{1,r-1} & \cdots & \binom{r}{l} \mu_{l,r-l} & \cdots & \binom{r}{r-1} \mu_{r-1,1} & \mu_{r,0} \end{array}$$

Lemma 1 (Pixel Intensity Reconstruction Property). *If the grid point locations $\{z_k\}_{k=1}^N$ are known and pairwise distinct, then the image I can be reconstructed from the Pascal Triangle $T^{N-1}(I)$ of order $N-1$. More specifically, knowledge of the entries of the right diagonal row of $T^{N-1}(I)$, i.e. $\{\mu_{j,0}\}_{j=0}^{N-1}$, is sufficient for image reconstruction.*

Proof. Recall the definition of the moments

$$\mu_{j,l} = \sum_{k=1}^N z_k^j \bar{z}_k^l \rho(z_k).$$

We consider the vector formed by the moments $\{\mu_{j,0}\}_{j=0}^{N-1}$, which can be written in matrix form as

$$\begin{pmatrix} \mu_{0,0} \\ \mu_{1,0} \\ \mu_{2,0} \\ \vdots \\ \mu_{N-1,0} \end{pmatrix} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ z_1 & z_2 & \cdots & z_N \\ z_1^2 & z_2^2 & \cdots & z_N^2 \\ \vdots & \vdots & \ddots & \vdots \\ z_1^{N-1} & z_2^{N-1} & \cdots & z_N^{N-1} \end{pmatrix} \begin{pmatrix} \rho(z_1) \\ \rho(z_2) \\ \vdots \\ \rho(z_N) \end{pmatrix}. \quad (2)$$

Observe that the coefficient matrix in (2) is a Vandermonde matrix. The Vandermonde matrix has full rank when $z_j \neq z_k$ for all distinct $j, k = 1, 2, \dots, N$. Thus, since the pixel locations are assumed to be distinct, we can reconstruct the pixel intensities $\{\rho(z_k)\}_{k=1}^N$ by inverting the coefficient matrix and multiplying by the moment vector on the left-hand-side. $\square$

Notice that if we consider the Pascal Triangle $T^N(I)$ of order N , then knowledge of the second right diagonal row of $T^N(I)$, i.e. $\{\mu_{j,1}\}_{j=0}^{N-1}$, is also sufficient for image reconstruction as long as the z_k 's are pairwise distinct and nonzero. This is because the vector formed by the moments $\{\mu_{j,1}\}_{j=0}^{N-1}$ can be written in matrix form as

$$\begin{pmatrix} \mu_{0,1} \\ \mu_{1,1} \\ \mu_{2,1} \\ \vdots \\ \mu_{N-1,1} \end{pmatrix} = \begin{pmatrix} \bar{z}_1 & \bar{z}_2 & \cdots & \bar{z}_N \\ z_1 \bar{z}_1 & z_2 \bar{z}_2 & \cdots & z_N \bar{z}_N \\ z_1^2 \bar{z}_1 & z_2^2 \bar{z}_2 & \cdots & z_N^2 \bar{z}_N \\ \vdots & \vdots & \ddots & \vdots \\ z_1^{N-1} \bar{z}_1 & z_2^{N-1} \bar{z}_2 & \cdots & z_N^{N-1} \bar{z}_N \end{pmatrix} \begin{pmatrix} \rho(z_1) \\ \rho(z_2) \\ \vdots \\ \rho(z_N) \end{pmatrix}, \quad (3)$$

$$= \begin{pmatrix} 1 & 1 & \cdots & 1 \\ z_1 & z_2 & \cdots & z_N \\ z_1^2 & z_2^2 & \cdots & z_N^2 \\ \vdots & \vdots & \ddots & \vdots \\ z_1^{N-1} & z_2^{N-1} & \cdots & z_N^{N-1} \end{pmatrix} \begin{pmatrix} \bar{z}_1 & 0 & \cdots & 0 \\ 0 & \bar{z}_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \bar{z}_N \end{pmatrix} \begin{pmatrix} \rho(z_1) \\ \rho(z_2) \\ \vdots \\ \rho(z_N) \end{pmatrix}. \quad (4)$$

The coefficient matrix in (3) is a Vandermonde matrix multiplied by a diagonal matrix. Assuming that the pixel locations are pairwise distinct insures that the Vandermonde matrix is invertible, and further assuming that they are nonzero insures invertibility of the diagonal matrix. Hence the coefficient matrix in (3) is nonsingular and we can reconstruct the pixel intensities $\{\rho(z_k)\}_{k=1}^N$ from $T^N(I)$ by inverting this coefficient matrix and multiplying by the

moment vector on the left-hand-side.

A similar argument can be used to show that, for any fixed l , the pixel intensities can be reconstructed from the moment vector $\{\mu_{j,l}\}_{j=0}^{N-1}$, which can be obtained from the Pascal Triangle $T^{N+l-1}(I)$ of order $N + l - 1$.

Remark: In practice, when reconstructing the pixel intensities of an image I , floating point errors in the matrix inversion can result in inaccuracies in the reconstructed image. In fact, the recovered pixel intensities may be complex valued. While the imaginary part of the result tends to be quite small, it is advantageous to first reformulate the problem to guarantee a real solution. One way to force the solution to be real is to separate Equation (2) into two sets of equations with real coefficients. More specifically, we can separate the equation system into its real part and its imaginary part, and combine these two real equation systems into one. After this, a real solution for the new equation system can be found, for example, by Singular Value Decomposition (SVD).

Lemma 2. ² *Given the moments matrix $\tau_N(I)$ of a discrete image I and an upper bound on the number N of pixels, one can reconstruct the pixel location z_k and the intensity $\rho(z_k)$ for all z_k such that $\rho(z_k) \neq 0$.*

Proof. If the number of pixels in the image I is strictly less than N , we can extend I to an image with N pixels by adding zero intensity pixels. Without loss of generality, we assume that $\rho(z_k) \neq 0$ for $k = 1, \dots, s$ and $\rho(z_k) = 0$ for $k = s + 1, \dots, N$. Consider the polynomial

$$P(t) = \prod_{k=1}^s (t - z_k) = t^s + \sum_{j=1}^s c_j t^{s-j}, \quad (5)$$

where the coefficients c_j are polynomials in the z_k 's.

Observe that $P(z_k) = 0$, $\forall k = 1, 2, \dots, s$. Therefore, we also have $\rho(z_k) \bar{z}_k^l P(z_k) = 0$, for any $l = 0, \dots, s - 1$. Summing all these equations over k 's, we get

²This result generalizes Proposition 1 in [1], which states that the vertices of a polygon are uniquely determined by a finite number of moments.

$$\begin{aligned}
& \sum_{k=1}^s \rho(z_k) \bar{z}_k^l P(z_k) = 0, \\
& \implies \sum_{k=1}^s \rho(z_k) \bar{z}_k^l (z_k^s + \sum_{j=1}^s c_j z_k^{s-j}) = 0, \\
& \implies \sum_{k=1}^s \rho(z_k) \bar{z}_k^l z_k^s + \sum_{j=1}^s c_j \sum_{k=1}^s \rho(z_k) \bar{z}_k^l z_k^{s-j} = 0, \\
& \implies \mu_{s,l} + \sum_{j=1}^s c_j \mu_{s-j,l} = 0, \\
& \implies \sum_{j=1}^s c_j \mu_{s-j,l} = -\mu_{s,l}, \quad l = 0, 1, \dots, s-1.
\end{aligned}$$

We write these last equations in matrix form:

$$\underbrace{\begin{pmatrix} \mu_{0,0} & \mu_{1,0} & \mu_{2,0} & \mu_{0,3} & \cdots & \mu_{s-1,0} \\ \mu_{0,1} & \mu_{1,1} & \mu_{2,1} & \cdots & \cdots & \mu_{s-1,1} \\ \mu_{0,2} & \mu_{1,2} & \cdots & \cdots & \cdots & \mu_{s-1,2} \\ \mu_{0,3} & \cdots & \cdots & \cdots & \cdots & \mu_{s-1,3} \\ \vdots & & & & & \vdots \\ \mu_{0,s-1} & \mu_{1,s-1} & \cdots & \cdots & \cdots & \mu_{s-1,s-1} \end{pmatrix}}_{\tau_s(I)} \begin{pmatrix} c_s \\ c_{s-1} \\ c_{s-2} \\ \vdots \\ c_1 \end{pmatrix} = - \begin{pmatrix} \mu_{s,0} \\ \mu_{s,1} \\ \mu_{s,2} \\ \vdots \\ \mu_{s,s-1} \end{pmatrix}. \quad (6)$$

From Equation (1) we know that

$$\tau_s(I) = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ z_1 & z_2 & \cdots & z_s \\ \vdots & \vdots & \ddots & \vdots \\ z_1^{s-1} & z_2^{s-1} & \cdots & z_s^{s-1} \end{pmatrix} \begin{pmatrix} \rho(z_1) & 0 & \cdots & 0 \\ 0 & \rho(z_2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \rho(z_s) \end{pmatrix} \begin{pmatrix} 1 & z_1 & z_1^2 & \cdots & z_1^{s-1} \\ 1 & z_2 & z_2^2 & \cdots & z_2^{s-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_s & z_s^2 & \cdots & z_s^{s-1} \end{pmatrix}.$$

Thus $\tau_s(I)$ is invertible, since the locations z_k are pairwise distinct and the pixel intensities $\rho(z_k)$ are nonzero. Hence we can solve the above equation system for $(c_s, c_{s-1}, \dots, c_1)$ by inverting $\tau_s(I)$ and multiplying by the vector on the right-hand-side of Equation (6).

Since the c_k 's determine the polynomial $P(t)$, we can solve for the roots of $P(t) = 0$, which are actually $\{z_k\}_{k=1}^s$. By Lemma 1, we can subsequently obtain the pixel intensities $\{\rho(z_k)\}_{k=1}^s$. $\square$

Remark: To determine the number of nonzero pixels, we can look at the rank of $\tau_N(I)$. Since $\tau_N(I) = Z^\dagger W Z$ by Equation (1) and $\text{rank}(Z^\dagger) = \text{rank}(Z) = N$, $\text{rank}(W) = s$, we can

conclude that $\text{rank}(\tau_N(I)) = s$.

Since the Pascal Triangle $T^{2N-2}(I)$ of the image I contains all the information needed to recover $\tau_N(I)$, we have the following corollary:

Corollary 1 (Image Reconstruction Property). *Given the Pascal Triangle $T^{2N-2}(I)$ of a discrete image I , one can reconstruct both the grid point locations $\{z_k\}_{k=1}^N$ and the corresponding intensities $\{\rho(z_k)\}_{k=1}^N$ for all those z_k such that $\rho(z_k) \neq 0$.*

2 Relationship with the Radon Transform

The Radon transform $f_\theta(r)$ is the projection of the image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ onto the straight line through the origin with direction vector $(\cos(\theta) \ \sin(\theta))^T$, i.e.

$$f_\theta(r) = \sum_{k \in S} \rho(z_k),$$

where $S = \{k | x_k \cos(\theta) + y_k \sin(\theta) = r, \ k = 1, 2, \dots, N\}$. Since $f_\theta(r)$ is a periodic function of θ with period 2π , any of its n th order moment $m_n(\theta)$ is also periodic with period 2π . It turns out that, for any $n = 0, 1, 2, \dots$, the coefficients of the Fourier series of $m_n(\theta)$ are given by the coefficients of row $(n+1)$ of $T^r(I)$ with $r \geq n$.

Lemma 3. *The n th order moment $m_n(\theta)$ of the radon transform $f_\theta(r)$ is given by the following linear combination of the $(n+1)$ th row entries of the Pascal Triangle $T^r(I)$ with $r \geq n$:*

$$m_n(\theta) = \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \mu_{l, n-l} e^{i(n-2l)\theta}. \quad (7)$$

Proof. For $n = 0$, we have

$$m_0(\theta) = \sum_{k=1}^N \rho(z_k) = \sum_{k=1}^N \rho(z_k) z_k^0 \bar{z}_k^0 = \mu_{0,0}.$$

For $n > 0$, we have

$$m_n(\theta) = \sum_r r^n f_\theta(r) = \sum_{k=1}^N (r_k(\theta))^n \rho(z_k),$$

where $r_k(\theta)$ is the projection of the vector $(x_k, y_k)^T$ onto the axis with angle $\theta \in (-\pi, \pi]$ with respect to x-axis. More precisely,

$$r_k(\theta) = x_k \cos \theta + y_k \sin \theta = \frac{1}{2}(z_k e^{-i\theta} + \bar{z}_k e^{i\theta}), \quad \forall k = 1, 2, \dots, N,$$

and therefore

$$\begin{aligned}
m_n(\theta) &= \sum_{k=1}^N (r_k(\theta))^n \rho(z_k), \\
&= \sum_{k=1}^N \left(\frac{1}{2} (z_k e^{-i\theta} + \bar{z}_k e^{i\theta}) \right)^n \rho(z_k), \\
&= \frac{1}{2^n} \sum_{k=1}^N (z_k e^{-i\theta} + \bar{z}_k e^{i\theta})^n \rho(z_k), \\
&= \frac{1}{2^n} \sum_{k=1}^N \left(\sum_{l=0}^n \binom{n}{l} z_k^l e^{-il\theta} \bar{z}_k^{n-l} e^{i(n-l)\theta} \right) \rho(z_k), \\
&= \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \left(\sum_{k=1}^N z_k^l \bar{z}_k^{n-l} \rho(z_k) \right) e^{-il\theta} e^{i(n-l)\theta}, \\
&= \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \mu_{l, n-l} e^{i(n-2l)\theta}.
\end{aligned}$$

□

Figure 1 and Figure 2 summarize the relationship between the Pascal Triangle and the Radon transform of an image when the pixel locations are known and unknown, respectively. Observe that a smaller number of rows of the Pascal Triangle are needed in order to reconstruct the image than if the pixel locations were known.




Figure 1: Relationship between the Pascal Triangle and the Radon transform of an image under the assumption that the pixel locations are unknown a priori.

3 Image Reconstruction from Samples

Lemma 4. *The last row of $T^n(I)$ can be reconstructed from $n + 1$ generic moment samples $m_n(\theta_1), m_n(\theta_2), \dots, m_n(\theta_{n+1})$.*




Figure 2: **Relationship between the Pascal Triangle and the Radon transform of an image under the assumption that the pixel locations z_k are fixed and known a priori.**

Proof. We first write Equation (7) in matrix form:

$$\begin{pmatrix} m_n(\theta_1) \\ m_n(\theta_2) \\ \vdots \\ m_n(\theta_{n+1}) \end{pmatrix} = \frac{1}{2^n} \begin{pmatrix} e^{in\theta_1} & \dots & e^{i(n-2l)\theta_1} & \dots & e^{-in\theta_1} \\ e^{in\theta_2} & \dots & e^{i(n-2l)\theta_2} & \dots & e^{-in\theta_2} \\ \vdots & \dots & \vdots & \dots & \vdots \\ e^{in\theta_{n+1}} & \dots & e^{i(n-2l)\theta_{n+1}} & \dots & e^{-in\theta_{n+1}} \end{pmatrix} \begin{pmatrix} \mu_{0,n} \\ \vdots \\ \binom{n}{l} \mu_{l,n-l} \\ \vdots \\ \mu_{n,0} \end{pmatrix}.$$

There is a unique solution for $(\mu_{0,n}, \dots, \binom{n}{l} \mu_{l,n-l}, \dots, \mu_{n,0})$ if and only if

$$\begin{aligned} & \det \left[\begin{pmatrix} e^{in\theta_1} & \dots & e^{i(n-2l)\theta_1} & \dots & e^{-in\theta_1} \\ e^{in\theta_2} & \dots & e^{i(n-2l)\theta_2} & \dots & e^{-in\theta_2} \\ \vdots & \dots & \vdots & \dots & \vdots \\ e^{in\theta_{n+1}} & \dots & e^{i(n-2l)\theta_{n+1}} & \dots & e^{-in\theta_{n+1}} \end{pmatrix} \right] \neq 0, \\ \iff & \det \left[\begin{pmatrix} e^{-in\theta_1} & 0 & 0 & \dots & 0 \\ 0 & e^{-in\theta_2} & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & \dots & e^{-in\theta_{n+1}} \end{pmatrix} \begin{pmatrix} e^{i2n\theta_1} & \dots & e^{i4\theta_1} & e^{i2\theta_1} & 1 \\ e^{i2n\theta_2} & \dots & e^{i4\theta_2} & e^{i2\theta_2} & 1 \\ \vdots & \dots & \vdots & \vdots & \vdots \\ e^{i2n\theta_{n+1}} & \dots & e^{i4\theta_{n+1}} & e^{i2\theta_{n+1}} & 1 \end{pmatrix} \right] \neq 0, \\ \iff & \det \left[\begin{pmatrix} e^{i2n\theta_1} & \dots & e^{i4\theta_1} & e^{i2\theta_1} & 1 \\ e^{i2n\theta_2} & \dots & e^{i4\theta_2} & e^{i2\theta_2} & 1 \\ \vdots & \dots & \vdots & \vdots & \vdots \\ e^{i2n\theta_{n+1}} & \dots & e^{i4\theta_{n+1}} & e^{i2\theta_{n+1}} & 1 \end{pmatrix} \right] \neq 0, \text{ since } e^{-in\theta_j} \neq 0, \forall j = 1, \dots, n+1. \end{aligned}$$

Observe that the above determinant is a Vandermonde determinant. It is nonzero if $e^{i2\theta_j} \neq e^{i2\theta_k}$ for all distinct $j, k = 1, \dots, n+1$. Therefore, if the θ_j 's are such that $e^{i2\theta_j} \neq e^{i2\theta_k}$

(thus the need to pick a generic sample set), we will get a unique solution for $(\mu_{0,n}, \dots, \binom{n}{l} \mu_{l,n-l}, \dots, \mu_{n,0})$. Hence we can reconstruct the last row of $T^n(I)$. $\square$

Corollary 2.³ *Given the grid point locations $\{z_k\}_{k=1}^N$, we can reconstruct the discrete image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ from the radon transform $f_{\theta_1}(r), f_{\theta_2}(r), \dots, f_{\theta_N}(r)$ at N fixed generic angles $\theta_1, \dots, \theta_N$.*

Proof. From the given radon transform of the image at different angles, we can calculate the moments $\{m_n(\theta_k) | n = 0, \dots, N-1, k = 1, \dots, n+1\}$. The conclusion followed by combining Lemma 1 and Lemma 3. $\square$

The diagram of Figure 3 thus commutes. Note that, one could use a similar argument along with Corollary 1 to show that $(2N-1)$ generic observations of $\{m_n(\theta_j), j = 1, \dots, n+1\}_{n=0}^{2N-2}$ would be needed to fully reconstruct the image $\{(z_k, \rho(z_k))\}_{k=1}^N$ with pixel positions z_k unknown.




Figure 3: Relationship between the Pascal Triangle $T^{N-1}(I)$ and the Radon transform.

4 Prolongation of Group Actions on the Moments and Invariantization

Let $(G, \cdot)$ be a group acting on the complex plane:

$$\begin{aligned} \cdot : G \times \mathbb{C} &\longrightarrow \mathbb{C} \\ (g, z) &= g \cdot z, \quad \forall g \in G, z \in \mathbb{C}. \end{aligned}$$

³This result generalizes Theorem 5.1 in [2], which states that a quadrature domain can be uniquely reconstructed by the line integral projections at finite angles.

This induces a group transformation $(G, \circ)$ of the discrete image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$, namely

$$g \circ \{(z_k, \rho(z_k))\}_{k=1}^N = \{(g \cdot z_k, \rho(z_k))\}_{k=1}^N, \quad \forall g \in G.$$

Then the induced transformation $(G, *)$ on moments $\{\mu_{j,l}\}_{j,l \in \mathbb{Z}_{\geq 0}}$ is

$$g * \{\mu_{j,l}\}_{j,l \in \mathbb{Z}_{\geq 0}} = g * \left\{ \sum_{k=1}^N z_k^j \bar{z}_k^l \rho(z_k) \right\}_{j,l \in \mathbb{Z}_{\geq 0}} = \left\{ \sum_{k=1}^N (g \cdot z_k)^j (\overline{g \cdot z_k})^l \rho(z_k) \right\}_{j,l \in \mathbb{Z}_{\geq 0}}.$$

In other words, the transformed moments are the moments of the transformed image.

Example 1: Consider the action of $G = \mathbb{C}$ on $\mathbb{C}$ by translation

$$(z_0, z) \mapsto z + z_0, \quad \forall z_0 \in G, \forall z \in \mathbb{C}.$$

Then the induced transformation on the image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ is

$$z_0 \circ \{(z_k, \rho(z_k))\}_{k=1}^N = \{(z_k + z_0, \rho(z_k))\}_{k=1}^N, \quad \forall z_0 \in G. \quad (8)$$

In other words, the image is translated horizontally with distance $x_0 = \text{Re}(z_0)$ and vertically with $y_0 = \text{Im}(z_0)$. The transformed complex moments are

$$\begin{aligned} \tilde{\mu}_{j,l} &= \sum_{k=1}^N \tilde{z}_k^j \bar{\tilde{z}}_k^l \rho(\tilde{z}_k) = \sum_{k=1}^N (z_k + z_0)^j (\bar{z}_k + \bar{z}_0)^l \rho(z_k), \\ &= \sum_{k=1}^N \left(\sum_{s=0}^j \binom{j}{s} z_k^s z_0^{j-s} \right) \left(\sum_{t=0}^l \binom{l}{t} \bar{z}_k^t \bar{z}_0^{l-t} \right) \rho(z_k), \\ &= \sum_{s=0}^j \sum_{t=0}^l \left(\sum_{k=1}^N z_k^s \bar{z}_k^t \rho(z_k) \right) \binom{j}{s} \binom{l}{t} z_0^{j-s} \bar{z}_0^{l-t}, \\ &= \sum_{s=0}^j \sum_{t=0}^l \mu_{s,t} \binom{j}{s} \binom{l}{t} z_0^{j-s} \bar{z}_0^{l-t}, \quad \forall j, l \in \mathbb{Z}_{\geq 0}. \end{aligned} \quad (9)$$

Written in matrix form, the transformation of the moment matrix $\tau_N(I)$ is

$$\tau_N(\tilde{I}) = A^\dagger \tau_N(I) A,$$

where $A = (a_{j,l})_{N \times N}$ is an upper-triangular matrix with $a_{j,l} = \binom{l-1}{j-1} z_0^{l-j}$, and $A^\dagger$ is the conjugate transpose of A .

Having obtained an explicit formula for the action of G on the moments, we follow Fels and Olver's moving frame method [3] to obtain a set of invariant functions of the moments. More

specifically, we consider the cross-section defined by $\tilde{\mu}_{1,0} = 0$. The group transformation that maps $\tau_N(I)$ to the cross-section is the moving frame $z_0 = -\frac{\mu_{1,0}}{\mu_{0,0}}$. By applying the moving frame to the moment matrix, we obtain the matrix

$$\tilde{\tau}_N(I) = A_0^\dagger \tau_N(I) A_0,$$

where $A_0 = \left(\binom{l-1}{j-1} \left(-\frac{\mu_{1,0}}{\mu_{0,0}}\right)^{l-j} \right)_{N \times N}$. By equivariance of the moving frame, all the entries of $\tilde{\tau}_N(I)$ are invariant under translation. One can check that theses entries $\tilde{\mu}_{j,l}$ are actually the centralized moments

$$\tilde{\mu}_{j,l} = \sum_{k=1}^N \left(z_k - \frac{\mu_{1,0}}{\mu_{0,0}}\right)^j \left(\bar{z}_k - \frac{\bar{\mu}_{1,0}}{\mu_{0,0}}\right)^l \rho(z_k), \quad j, l \in \mathbb{Z}_{\geq 0}. \quad (10)$$

By normalizing (i.e. applying the moving frame transformation to) the coordinates of $T^r(I)$, we obtain the translation invariant Pascal Triangle $T_{trans}^r(I)$ for a discrete image I :

$$\begin{array}{ccccccc} & & & & & & \tilde{\mu}_{0,0} \\ & & & & & & \tilde{\mu}_{0,1} & \tilde{\mu}_{1,0} \\ & & & & & & \tilde{\mu}_{0,2} & 2\tilde{\mu}_{1,1} & \tilde{\mu}_{2,0} \\ & & & & & & \tilde{\mu}_{0,3} & 3\tilde{\mu}_{1,2} & 3\tilde{\mu}_{2,1} & \tilde{\mu}_{3,0} \\ & & & & & & \tilde{\mu}_{0,4} & 4\tilde{\mu}_{1,3} & 6\tilde{\mu}_{2,2} & 4\tilde{\mu}_{3,1} & \tilde{\mu}_{4,0} \\ & & & & & & & & & & \vdots \\ & & & & & & \tilde{\mu}_{0,r} & \binom{r}{1} \tilde{\mu}_{1,r-1} & \cdots & \binom{r}{l} \tilde{\mu}_{l,r-l} & \cdots & \binom{r}{r-1} \tilde{\mu}_{r-1,1} & \tilde{\mu}_{r,0} \end{array}$$

Observe that the corresponding n -th order central moment $\tilde{m}_n(\theta)$ of the image,

$$\tilde{m}_n(\theta) = \sum_{k=1}^N \left(r_k(\theta) - r_0(\theta)\right)^n \rho(x_k, y_k),$$

where $r_0(\theta) = x_0 \cos \theta + y_0 \sin \theta$ is the projection of the centroid, is invariant under translations.

Lemma 5 (Orbit Separation Property of $T_{trans}^{2N-2}(I)$). *Let I_1, I_2 be two discrete gray scale images with the same number N of pixels. There exists a translation $g \in \mathbb{C}$ such that $g \circ I_1 = I_2$, where $\circ$ is defined as in (8) $\iff T_{trans}^r(I_1) = T_{trans}^r(I_2)$ for $r \geq 2N - 2$.*

Proof. $\Rightarrow$ If $\exists g \in G$ such that $g \circ I_1 = I_2$, we have $z_k^{(2)} = z_k^{(1)} + z_0$ and $\rho_2(z_k^{(2)}) = \rho_1(z_k^{(1)})$, for some $z_0 \in \mathbb{C}$, $k = 1, \dots, N$. From Equation (9) we know that

$$\mu_{0,0}^{(2)} = \mu_{0,0}^{(1)}, \quad \mu_{1,0}^{(2)} = \mu_{0,0}^{(1)} z_0 + \mu_{1,0}^{(1)}.$$

Hence

$$\frac{\mu_{1,0}^{(2)}}{\mu_{0,0}^{(2)}} = \frac{\mu_{1,0}^{(1)}}{\mu_{0,0}^{(1)}} + z_0.$$

Then applying Equation (10), we can get for any $j, l \in \mathbb{Z}_{\geq 0}$

$$\tilde{\mu}_{j,l}^{(1)} = \sum_{k=1}^N \left(z_k^{(1)} - \frac{\mu_{1,0}^{(1)}}{\mu_{0,0}^{(1)}} \right)^j \left(\bar{z}_k^{(1)} - \frac{\bar{\mu}_{1,0}^{(1)}}{\mu_{0,0}^{(1)}} \right)^l \rho_1(z_k^{(1)}) = \sum_{k=1}^N \left(z_k^{(2)} - \frac{\mu_{1,0}^{(1)}}{\mu_{0,0}^{(1)}} z_0 \right)^j \left(\bar{z}_k - \frac{\bar{\mu}_{1,0}^{(1)}}{\mu_{0,0}^{(1)}} \bar{z}_0 \right)^l \rho_2(z_k^{(2)}) = \tilde{\mu}_{j,l}^{(2)}.$$

Therefore $T_{trans}^r(I_1) = T_{trans}^r(I_2)$ for any $r \in \mathbb{Z}_{\geq 0}$.

$\Leftarrow$ If $T_{trans}^r(I_1) = T_{trans}^r(I_2)$ for $r \geq 2N-2$, from Corollary 1, we conclude that $I_1^{trans} = I_2^{trans}$, i.e.

$$\left\{ (z_{k,trans}^{(1)}, \rho_1(z_{k,trans}^{(1)})) \right\}_{k=1}^N = \left\{ (z_{k,trans}^{(2)}, \rho_2(z_{k,trans}^{(2)})) \right\}_{k=1}^N.$$

Hence $\exists z_0, z'_0 \in \mathbb{C}$ s.t. $z_{k,trans}^{(1)} = z_k^{(1)} + z_0$, $z_{k,trans}^{(2)} = z_k^{(2)} + z'_0$ with $\rho_1(z_{k,trans}^{(1)}) = \rho_1(z_k^{(1)})$ and $\rho_2(z_{k,trans}^{(2)}) = \rho_2(z_k^{(2)})$ for any $k = 1, 2, \dots, N$. Without loss of generality, we assume that $z_{k,trans}^{(1)} = z_{k,trans}^{(2)}$ and $\rho_1(z_{k,trans}^{(1)}) = \rho_2(z_{k,trans}^{(2)})$. Therefore $\exists z_0 - z'_0 \in \mathbb{C} = G$ satisfying

$$z_k^{(1)} + z_0 - z'_0 = z_k^{(2)}, \quad \rho_1(z_k^{(1)} + (z_0 - z'_0)) = \rho_1(z_k^{(1)}) = \rho_2(z_k^{(2)}),$$

i.e. $\exists g = z'_0 - z_0 \in G = \mathbb{C}$ such that $g \circ I_1 = I_2$. $\square$

Remark: Without loss of generality, we can assume that the two images have the same number of pixels by simply adding zero valued pixels to the smaller image.

Example 2: Consider the action of $G = \mathbb{R}_+$ on $\mathbb{C}$ by scaling

$$(\lambda, z) \mapsto \lambda z, \quad \forall \lambda \in G, \forall z \in \mathbb{C}.$$

Then the induced transformation on the image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ is

$$\lambda \circ \{(z_k, \rho(z_k))\}_{k=1}^N = \{(\lambda z_k, \rho(z_k))\}_{k=1}^N, \quad \forall \lambda \in G. \quad (11)$$

In other words, the image is scaled by a factor λ both horizontally and vertically. Then the transformed complex moments are

$$\begin{aligned}
\hat{\mu}_{j,l} &= \sum_{k=1}^N \hat{z}_k^j \bar{\hat{z}}_k^l \rho(\hat{z}_k) = \sum_{k=1}^N (\lambda z_k)^j (\lambda \bar{z}_k)^l \rho(z_k), \\
&= \sum_{k=1}^N \lambda^{j+l} z_k^j \bar{z}_k^l \rho(z_k) = \lambda^{j+l} \sum_{k=1}^N z_k^j \bar{z}_k^l \rho(z_k), \\
&= \mu_{j,l} \lambda^{j+l}, \quad \forall j, l \in \mathbb{Z}_{\geq 0}.
\end{aligned}$$

Written in matrix form, the moment matrix for the new image $\hat{I}$ after scaling is

$$\tau_N(\hat{I}) = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda & 0 & \cdots & 0 \\ 0 & 0 & \lambda^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & \lambda^{N-1} \end{pmatrix} \tau_N(I) \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda & 0 & \cdots & 0 \\ 0 & 0 & \lambda^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & \lambda^{N-1} \end{pmatrix}.$$

Again, we use the moving frame method of Fels and Olver to obtain a set of invariant functions of the moments. Notice that $\hat{\mu}_{1,1} = \sum_{k=1}^N \hat{z}_k \bar{\hat{z}}_k \rho(\hat{z}_k) = \sum_{k=1}^N (\hat{x}_k^2 + \hat{y}_k^2) \rho(\hat{z}_k) \neq 0$ unless all $\rho(z_k)$ are zero. We consider the cross-section defined by $\hat{\mu}_{1,1} = 1$. The group transformation that maps $\tau_N(I)$ to the cross-section is the moving frame $\lambda = (\mu_{1,1})^{-\frac{1}{2}}$. By applying the moving frame to the moment matrix, we obtain the matrix

$$\hat{\tau}_N(I) = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & (\mu_{1,1})^{-\frac{1}{2}} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & (\mu_{1,1})^{-\frac{N-1}{2}} \end{pmatrix} \tau_N(I) \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & (\mu_{1,1})^{-\frac{1}{2}} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & (\mu_{1,1})^{-\frac{N-1}{2}} \end{pmatrix}.$$

By equivariance of the moving frame, all the entries of $\hat{\tau}_N(I)$ are invariant under scaling.

By normalizing the coordinates of $T^r(I)$, we obtain the scaling invariant Pascal Triangle $T_{scale}^r(I)$ for a discrete image I :

$$\begin{array}{ccccccc}
& & & & \mu_{0,0} & & \\
& & & & \mu_{0,1}/\sqrt{\mu_{1,1}} & \mu_{1,0}/\sqrt{\mu_{1,1}} & \\
& & & \mu_{0,2}/\mu_{1,1} & 2 & \mu_{2,0}/\mu_{1,1} & \\
& & \mu_{0,3}/\mu_{1,1}^{\frac{3}{2}} & 3\mu_{1,2}/\mu_{1,1}^{\frac{3}{2}} & 3\mu_{2,1}/\mu_{1,1}^{\frac{3}{2}} & \mu_{3,0}/\mu_{1,1}^{\frac{3}{2}} & \\
& \mu_{0,4}/\mu_{1,1}^2 & 4\mu_{1,3}/\mu_{1,1}^2 & 6\mu_{2,2}/\mu_{1,1}^2 & 4\mu_{3,1}/\mu_{1,1}^2 & \mu_{4,0}/\mu_{1,1}^2 & \\
& & & \vdots & & & \\
\mu_{0,r}/\mu_{1,1}^{\frac{r}{2}} & \binom{r}{1} \mu_{1,r-1}/\mu_{1,1}^{\frac{r}{2}} & \cdots & \binom{r}{l} \mu_{l,r-l}/\mu_{1,1}^{\frac{r}{2}} & \cdots & \binom{r}{r-1} \mu_{r-1,1}/\mu_{1,1}^{\frac{r}{2}} & \mu_{r,0}/\mu_{1,1}^{\frac{r}{2}}
\end{array}$$

Observe that the corresponding n -th order normalized moment $\hat{m}_n(\theta) = m_n(\theta)/\mu_{1,1}^{\frac{n}{2}}$ is invariant under scaling.

Lemma 6 (Orbit Separation Property of $T_{scale}^{2N-2}(I)$). *Let I_1, I_2 be two discrete gray scale images with the same number N of pixels. There exists a scaling $g \in \mathbb{R}_+$ such that $g \circ I_1 = I_2$, where $\circ$ is defined as in (11) $\iff T_{scale}^r(I_1) = T_{scale}^r(I_2)$ for $r \geq 2N - 2$.*

Proof. $\Rightarrow$ If $\exists g \in G$ such that $g \circ I_1 = I_2$, we have $z_k^{(2)} = \lambda z_k^{(1)}$ and $\rho_1(z_k^{(1)}) = \rho_2(z_k^{(2)})$ for some $\lambda \in \mathbb{R}_+$, $k = 1, \dots, N$. Since for I_1 and I_2 , the corresponding scaling invariant moments are

$$\begin{aligned} \hat{\mu}_{j,l}^{(1)} &= \frac{\mu_{j,l}^{(1)}}{(\mu_{1,1}^{(1)})^{\frac{j+l}{2}}} = \frac{\sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})}{\left(\sum_{k=1}^N z_k^{(1)} \bar{z}_k^{(1)} \rho_1(z_k^{(1)})\right)^{\frac{j+l}{2}}}, \\ \hat{\mu}_{j,l}^{(2)} &= \frac{\mu_{j,l}^{(2)}}{(\mu_{1,1}^{(2)})^{\frac{j+l}{2}}} = \frac{\sum_{k=1}^N (z_k^{(2)})^j (\bar{z}_k^{(2)})^l \rho_2(z_k^{(2)})}{\left(\sum_{k=1}^N z_k^{(2)} \bar{z}_k^{(2)} \rho_2(z_k^{(2)})\right)^{\frac{j+l}{2}}} \\ &= \frac{\sum_{k=1}^N (\lambda z_k^{(1)})^j (\lambda \bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})}{\left(\sum_{k=1}^N (\lambda z_k^{(1)}) (\lambda \bar{z}_k^{(1)}) \rho_1(z_k^{(1)})\right)^{\frac{j+l}{2}}} = \frac{\sum_{k=1}^N \lambda^{j+l} (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})}{\left(\sum_{k=1}^N \lambda^2 z_k^{(1)} \bar{z}_k^{(1)} \rho_1(z_k^{(1)})\right)^{\frac{j+l}{2}}} \\ &= \frac{\lambda^{j+l} \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})}{\lambda^{j+l} \left(\sum_{k=1}^N z_k^{(1)} \bar{z}_k^{(1)} \rho_1(z_k^{(1)})\right)^{\frac{j+l}{2}}} = \frac{\sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})}{\left(\sum_{k=1}^N z_k^{(1)} \bar{z}_k^{(1)} \rho_1(z_k^{(1)})\right)^{\frac{j+l}{2}}} = \hat{\mu}_{j,l}^{(1)}. \end{aligned}$$

Therefore $T_{scale}^r(I_1) = T_{scale}^r(I_2)$ for any $r \in \mathbb{Z}_{\geq 0}$.

$\Leftarrow$ If $T_{scale}^r(I_1) = T_{scale}^r(I_2)$ for $r \geq 2N - 2$, from Corollary 1, we conclude that $I_1^{scale} = I_2^{scale}$, i.e.

$$\{(z_{k,scale}^{(1)}, \rho_1(z_{k,scale}^{(1)}))\}_{k=1}^N = \{(z_{k,scale}^{(2)}, \rho_2(z_{k,scale}^{(2)}))\}_{k=1}^N.$$

Hence $\exists \lambda_1, \lambda_2 \in \mathbb{R}_+$ s.t. $z_{k,scale}^{(1)} = \lambda_1 z_k^{(1)}$, $z_{k,scale}^{(2)} = \lambda_2 z_k^{(2)}$ with $\rho_1(z_{k,scale}^{(1)}) = \rho_1(z_k^{(1)})$ and $\rho_2(z_{k,scale}^{(2)}) = \rho_2(z_k^{(2)})$ for any $k = 1, 2, \dots, N$. After relabeling, we have $z_{k,scale}^{(1)} = z_k^{(1)}$ and $\rho_1(z_{k,scale}^{(1)}) = \rho_2(z_{k,scale}^{(2)})$. Then $\exists \frac{\lambda_1}{\lambda_2} \in \mathbb{R}_+ = G$ satisfying

$$z_k^{(1)} \frac{\lambda_1}{\lambda_2} = z_k^{(2)}, \quad \rho_1(z_k^{(1)} \frac{\lambda_2}{\lambda_1}) = \rho_1(z_k^{(1)}) = \rho_2(z_k^{(2)}),$$

i.e. $\exists g = \frac{\lambda_2}{\lambda_1} \in G = \mathbb{R}_+$ such that $g \circ I_1 = I_2$. □

More generally, consider the action of group G of diagonal matrices on $\mathbb{R}_+^2$ by scaling

$$\left(\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}\right) \mapsto \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \lambda_1 x \\ \lambda_2 y \end{pmatrix}, \quad \forall \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \in G, \forall \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2.$$

Then the induced transformation on the image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ is

$$\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \circ \{(z_k, \rho(z_k))\}_{k=1}^N = \{(\lambda_1 x_k + i\lambda_2 y_k, \rho(z_k))\}_{k=1}^N, \quad \forall \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \in G, z_k = x_k + iy_k.$$

In other words, the image is scaled by a factor λ_1 horizontally and scaled by λ_2 vertically.

Notice that after the transformation, the pixel coordinates become

$$\begin{aligned} \hat{z}_k &= \lambda_1 x_k + i\lambda_2 y_k = \lambda_1 \frac{z_k + \bar{z}_k}{2} + i\lambda_2 \frac{z_k - \bar{z}_k}{2i} = \frac{\lambda_1 + \lambda_2}{2} z_k + \frac{\lambda_1 - \lambda_2}{2} \bar{z}_k, \\ \bar{\hat{z}}_k &= \lambda_1 x_k - i\lambda_2 y_k = \lambda_1 \frac{z_k + \bar{z}_k}{2} - i\lambda_2 \frac{z_k - \bar{z}_k}{2i} = \frac{\lambda_1 - \lambda_2}{2} z_k + \frac{\lambda_1 + \lambda_2}{2} \bar{z}_k. \end{aligned}$$

Then we have the transformed complex moments

$$\begin{aligned} \hat{\mu}_{j,l} &= \sum_{k=1}^N \hat{z}_k^j \bar{\hat{z}}_k^l \rho(\hat{z}_k) = \sum_{k=1}^N \left(\frac{\lambda_1 + \lambda_2}{2} z_k + \frac{\lambda_1 - \lambda_2}{2} \bar{z}_k \right)^j \left(\frac{\lambda_1 - \lambda_2}{2} z_k + \frac{\lambda_1 + \lambda_2}{2} \bar{z}_k \right)^l \rho(z_k), \\ &= \sum_{k=1}^N \rho(z_k) \left[\sum_{s=0}^j \binom{j}{s} \left(\frac{\lambda_1 + \lambda_2}{2} \right)^s z_k^s \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{j-s} \bar{z}_k^{j-s} \right] \left[\sum_{t=0}^l \binom{l}{t} \left(\frac{\lambda_1 - \lambda_2}{2} \right)^t z_k^t \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{l-t} \bar{z}_k^{l-t} \right], \\ &= \sum_{k=1}^N \rho(z_k) \left[\sum_{s=0}^j \sum_{t=0}^l \binom{j}{s} \binom{l}{t} \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{l-t+s} \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{j-s+t} z_k^{s+t} \bar{z}_k^{j+l-s-t} \right], \\ &= \sum_{s=0}^j \sum_{t=0}^l \binom{j}{s} \binom{l}{t} \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{l-t+s} \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{j-s+t} \left[\sum_{k=1}^N \rho(z_k) z_k^{s+t} \bar{z}_k^{j+l-s-t} \right], \\ &= \frac{1}{2^{j+l}} \sum_{s=0}^j \sum_{t=0}^l \binom{j}{s} \binom{l}{t} (\lambda_1 + \lambda_2)^{l-t+s} (\lambda_1 - \lambda_2)^{j-s+t} \mu_{s+t, j+l-s-t}, \quad \forall j, l \in \mathbb{Z}_{\geq 0}, \end{aligned}$$

which is a linear combination of the last row of the Pascal Triangle $T^{j+l}(I)$.

Example 3: Consider the action of $G = \{z \in \mathbb{C} \mid |z| = 1\}$ on $\mathbb{C}$ by rotation

$$(e^{i\theta_0}, z) \mapsto ze^{i\theta_0}, \quad \forall e^{i\theta_0} \in G, \forall z \in \mathbb{C}.$$

Then the induced transformation on the image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$ is

$$e^{i\theta_0} \circ \{(z_k, \rho(z_k))\}_{k=1}^N = \{(e^{i\theta_0} z_k, \rho(z_k))\}_{k=1}^N, \quad \forall e^{i\theta_0} \in G. \quad (12)$$

In other words, the image is rotated counterclockwise with an angle θ_0 . The transformed

complex moments are

$$\begin{aligned}
\mu'_{j,l} &= \sum_{k=1}^N z_k'^j \bar{z}_k'^l \rho(z_k') = \sum_{k=1}^N (z_k e^{i\theta_0})^j (\bar{z}_k e^{-i\theta_0})^l \rho(z_k), \\
&= \sum_{k=1}^N e^{i(j-l)\theta_0} z_k^j \bar{z}_k^l \rho(z_k) = e^{i(j-l)\theta_0} \sum_{k=1}^N z_k^j \bar{z}_k^l \rho(z_k), \\
&= \mu_{j,l} e^{i(j-l)\theta_0}, \quad \forall j, l \in \mathbb{Z}_{\geq 0}.
\end{aligned}$$

Written in matrix form, the moment matrix for the new image I' after rotation is

$$\tau_N(I') = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{-i\theta_0} & 0 & \cdots & 0 \\ 0 & 0 & e^{-i2\theta_0} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{-i(N-1)\theta_0} \end{pmatrix} \tau_N(I) \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{i\theta_0} & 0 & \cdots & 0 \\ 0 & 0 & e^{i2\theta_0} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{i(N-1)\theta_0} \end{pmatrix}.$$

Now we will use the moving frame method of Fels and Olver to obtain a set of invariant functions of the moments. If $\mu_{0,2} \neq 0$, we normalize the imaginary part of $\mu'_{0,2}$ to zero by specifying the rotation angle θ_0 . Since $\mu'_{0,2} = \mu_{0,2} e^{-i2\theta_0}$, looking at $\vec{\mu}_{0,2}$ as a vector in $\mathbb{R}^2$ representing the complex number $\mu_{0,2}$, we set

$$2\theta_0 = \angle(\vec{\mu}_{0,2}, \vec{e}_1) + 2k\pi, \quad k \in \mathbb{Z}.$$

Here $\angle(\vec{x}, \vec{y}) = \tan^{-1}(\frac{y_2}{y_1}) - \tan^{-1}(\frac{x_2}{x_1})$ denotes the angle from $\vec{x}$ to $\vec{y}$, $\vec{e}_1 = (1, 0)^T$ is one of the standard basis of $\mathbb{R}^2$. The real part of $\mu'_{0,2}$ then reduces to its magnitude $|\mu_{0,2}|$.

Since $\theta_0 \in (-\pi, \pi]$, $2\theta_0 \in (-2\pi, 2\pi]$. To uniquely determine the value of θ_0 , we consider $\mu'_{1,2} = \mu_{1,2} e^{-i\theta_0}$. We choose θ_0 such that $\text{Re}(\mu'_{1,2}) \geq 0$, which leads to the moving frame formulae

$$\theta_0 = \begin{cases} \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) & \text{if } \angle(\vec{\mu}_{1,2}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) \in [-\frac{\pi}{2}, \frac{\pi}{2}], \\ \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) + \pi & \text{if } \angle(\vec{\mu}_{1,2}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) \in (\frac{\pi}{2}, \frac{3\pi}{2}], \\ \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) - \pi & \text{if } \angle(\vec{\mu}_{1,2}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}, \vec{e}_1) \in [-\frac{3\pi}{2}, \frac{\pi}{2}). \end{cases} \quad (13)$$

By applying the moving frame to the moment matrix, we obtain the matrix

$$\tau'_N(I) = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{-i\theta_0} & 0 & \cdots & 0 \\ 0 & 0 & e^{-i2\theta_0} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{-i(N-1)\theta_0} \end{pmatrix} \tau_N(I) \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{i\theta_0} & 0 & \cdots & 0 \\ 0 & 0 & e^{i2\theta_0} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{i(N-1)\theta_0} \end{pmatrix},$$

with θ_0 satisfying (13). By equivariance of the moving frame, all the entries of $\tau'_N(I)$ are invariant under rotation.

By normalizing the coordinates of $T^r(I)$, we obtain the rotational invariant Pascal Triangle $T_{rotate}^r(I)$ for a discrete image I :

$$\begin{array}{ccccccc}
& & & & \mu_{0,0} & & \\
& & & & \mu_{0,1}e^{-i\theta_0} & \mu_{1,0}e^{i\theta_0} & \\
& & & |\mu_{0,2}| & 2\mu_{1,1} & |\mu_{2,0}| & \\
& & \mu_{0,3}e^{-i3\theta_0} & 3\mu_{1,2}e^{-i\theta_0} & 3\mu_{2,1}e^{i\theta_0} & \mu_{3,0}e^{i3\theta_0} & \\
\mu_{0,4}e^{-i4\theta_0} & 4\mu_{1,3}e^{-i2\theta_0} & 6\mu_{2,2} & 4\mu_{3,1}e^{i2\theta_0} & \mu_{4,0}e^{i4\theta_0} & & \\
& & & & \vdots & & \\
\mu_{0,r}e^{-ir\theta_0} & \binom{r}{1}\mu_{1,r-1}e^{-i(r-2)\theta_0} & \dots & \binom{r}{l}\mu_{l,r-l}e^{-i(r-2l)\theta_0} & \dots & \binom{r}{r-1}\mu_{r-1,1}e^{i(r-2)\theta_0} & \mu_{r,0}e^{ir\theta_0}
\end{array}$$

Observe that the corresponding n -th order moment $m'_n(\theta) = m_n(\theta - \theta_0)$, with θ_0 defined as in (13), is invariant under rotations.

Lemma 7 (Orbit Separation Property of $T_{rotate}^{2N-2}(I)$). *Let I_1, I_2 be two discrete gray scale images with the same number N of pixels. There exists a rotation $g \in \{z \in \mathbb{C} \mid |z| = 1\}$ such that $g \circ I_1 = I_2$, where $\circ$ is defined as in (12) $\iff T_{rotate}^r(I_1) = T_{rotate}^r(I_2)$ for $r \geq 2N - 2$.*

Proof. $\Rightarrow$ If $\exists g \in G$ such that $g \circ I_1 = I_2$, we have $z_k^{(2)} = z_k^{(1)}e^{i\theta_0}$ and $\rho_1(z_k^{(1)}) = \rho_2(z_k^{(2)})$, for some $\theta_0 \in \{z \in \mathbb{C} \mid |z| = 1\}$, $k = 1, \dots, N$. For I_1 and I_2 , the corresponding scaling invariant moments are

$$\begin{aligned}
\hat{\mu}_{j,l}^{(1)} &= \mu_{j,l}^{(1)}e^{-i(l-j)\theta_1} = \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})e^{i(j-l)\theta_1}, \\
\hat{\mu}_{j,l}^{(2)} &= \mu_{j,l}^{(2)}e^{-i(l-j)\theta_2} = \sum_{k=1}^N (z_k^{(2)})^j (\bar{z}_k^{(2)})^l \rho_2(z_k^{(2)})e^{i(j-l)\theta_2} \\
&= \sum_{k=1}^N (z_k^{(1)}e^{i\theta_0})^j (\bar{z}_k^{(1)}e^{-i\theta_0})^l \rho_1(z_k^{(1)})e^{i(j-l)\theta_2} \\
&= \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)})e^{i(j-l)(\theta_0+\theta_2)}.
\end{aligned}$$

Since $\mu_{0,2}^{(2)} = \mu_{0,2}^{(1)}e^{-i2\theta_0}$, $\mu_{1,2}^{(2)} = \mu_{1,2}^{(1)}e^{-i\theta_0}$, we have

$$\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) = \angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) - 2\theta_0 + 2k_1\pi, \quad \angle(\vec{\mu}_{1,2}^{(2)}, \vec{e}_1) = \angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \theta_0 + 2k_2\pi, \quad k_1, k_2 \in \mathbb{Z}. \quad (14)$$

Notice that $\angle(\vec{\mu}_{0,2}^{(l)}, \vec{e}_1), \angle(\vec{\mu}_{1,2}^{(l)}, \vec{e}_1), \theta_0 \in (-\pi, \pi]$, for $l = 1, 2$. Hence $k_1, k_2 = 0, \pm 1$. To decide θ_1 and θ_2 , we consider

$$\angle(\vec{\mu}_{1,2}^{(2)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) = \angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) + (2k_2 - k_1)\pi \quad (15)$$

by using (14).

If $\angle(\vec{\mu}_{1,2}^{(2)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, from (13) we know that $\theta_2 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1)$. Since $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{3\pi}{2}, \frac{3\pi}{2}]$, then $2k_2 - k_1$ is either 0 or ± 1 in (15). If $k_1 = 0, k_2 = 0$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1)$. Then from (14) we have $\theta_2 = \theta_1 - \theta_0$. Thus $\hat{\mu}_{j,l}^{(2)} = \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)}) e^{i(j-l)\theta_1} = \hat{\mu}_{j,l}^{(1)}$. If $k_1 = 1, k_2 = 0$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in (\frac{\pi}{2}, \frac{3\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) + \pi$. Then from (14) we still have $\theta_2 = \theta_1 - \theta_0$. Thus $\hat{\mu}_{j,l}^{(2)} = \hat{\mu}_{j,l}^{(1)}$. If $k_1 = 1, k_2 = 1$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{3\pi}{2}, -\frac{\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) - \pi$. Then from (14) we have $\theta_2 = \theta_1 - \theta_0 + 2\pi$. Thus $\hat{\mu}_{j,l}^{(2)} = \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)}) e^{i(j-l)\theta_1} = \hat{\mu}_{j,l}^{(1)}$.

If $\angle(\vec{\mu}_{1,2}^{(2)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) \in (\frac{\pi}{2}, \frac{3\pi}{2}]$, from (13) we know that $\theta_2 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) + \pi$. Since $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{3\pi}{2}, \frac{3\pi}{2}]$, then $2k_2 - k_1 = 0, 1, 2$ in (15). If $k_1 = 1, k_2 = 1$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1)$. Then from (14) we have $\theta_2 - \pi = \theta_1 - \theta_0 + \pi$. Thus $\hat{\mu}_{j,l}^{(2)} = \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)}) e^{i(j-l)\theta_1} = \hat{\mu}_{j,l}^{(1)}$. If $k_1 = 0, k_2 = 0$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in (\frac{\pi}{2}, \frac{3\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) + \pi$. Then from (14) we have $\theta_2 - \pi = \theta_1 - \theta_0 - \pi$. Thus $\hat{\mu}_{j,l}^{(2)} = \hat{\mu}_{j,l}^{(1)}$. If $k_1 = 0, k_2 = 1$, there is $\angle(\vec{\mu}_{1,2}^{(1)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) \in [-\frac{3\pi}{2}, -\frac{\pi}{2}]$. Hence $\theta_1 = \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(1)}, \vec{e}_1) - \pi$. Then from (14) we have $\theta_2 - \pi = \theta_1 - \theta_0 + \pi$. Thus $\hat{\mu}_{j,l}^{(2)} = \sum_{k=1}^N (z_k^{(1)})^j (\bar{z}_k^{(1)})^l \rho_1(z_k^{(1)}) e^{i(j-l)\theta_1} = \hat{\mu}_{j,l}^{(1)}$.

If $\angle(\vec{\mu}_{1,2}^{(2)}, \vec{e}_1) - \frac{1}{2}\angle(\vec{\mu}_{0,2}^{(2)}, \vec{e}_1) \in [-\frac{3\pi}{2}, -\frac{\pi}{2}]$, through the similar discussion, we can still conclude that $\hat{\mu}_{j,l}^{(2)} = \hat{\mu}_{j,l}^{(1)}$. Therefore $T_{scale}^r(I_1) = T_{scale}^r(I_2)$ for any $r \in \mathbb{Z}_{\geq 0}$.

$\Leftarrow$ If $T_{rotate}^r(I_1) = T_{rotate}^r(I_2)$ for $r \geq 2N - 2$, from Corollary 1, we conclude that $I_1^{rotate} = I_2^{rotate}$, i.e.

$$\{(z_{k,rotate}^{(1)}, \rho(z_{k,rotate}^{(1)}))\}_{k=1}^N = \{(z_{k,rotate}^{(2)}, \rho(z_{k,rotate}^{(2)}))\}_{k=1}^N.$$

Hence $\exists \theta_1, \theta_2 \in \{z \in \mathbb{C} \mid |z| = 1\}$ s.t. $z_{k,rotate}^{(1)} = z_k^{(1)} e^{i\theta_1}$, $z_{k,rotate}^{(2)} = z_k^{(2)} e^{i\theta_2}$ with $\rho_1(z_{k,rotate}^{(1)}) = \rho_1(z_k^{(1)})$ and $\rho_2(z_{k,rotate}^{(2)}) = \rho_2(z_k^{(2)})$ for any $k = 1, 2, \dots, N$. Without loss of generality, we assume $z_{k,rotate}^{(1)} = z_k^{(2)}$ and $\rho_1(z_{k,rotate}^{(1)}) = \rho_2(z_k^{(2)})$. Then $\exists \theta_1 - \theta_2 \in \{z \in \mathbb{C} \mid |z| = 1\} = G$ satisfying

$$z_k^{(1)} e^{i(\theta_1 - \theta_2)} = z_k^{(2)}, \quad \rho_1(z_k^{(1)} e^{i(\theta_1 - \theta_2)}) = \rho_1(z_k^{(1)}) = \rho_2(z_k^{(2)}),$$

i.e. $\exists g = e^{i(\theta_1 - \theta_2)} \in G = \{z \in \mathbb{C} \mid |z| = 1\}$ such that $g \circ I_1 = I_2$. $\square$

In conclusion, we have the translation, scaling and rotation invariant Pascal Triangle $T_{int}^r(I)$:

$$\begin{array}{ccccccc}
& & & & \tilde{\mu}_{0,0} & & \\
& & & & \tilde{\mu}_{0,1}e^{-i\theta_0}/\sqrt{\tilde{\mu}_{1,1}} & \tilde{\mu}_{1,0}e^{i\theta_0}/\sqrt{\tilde{\mu}_{1,1}} & \\
& & & |\tilde{\mu}_{0,2}|/\tilde{\mu}_{1,1} & 2\tilde{\mu}_{1,1}/\tilde{\mu}_{1,1} & |\tilde{\mu}_{2,0}|/\tilde{\mu}_{1,1} & \\
& & \tilde{\mu}_{0,3}e^{-i3\theta_0}/\tilde{\mu}_{1,1}^{\frac{3}{2}} & 3\tilde{\mu}_{1,2}e^{-i\theta_0}/\tilde{\mu}_{1,1}^{\frac{3}{2}} & 3\tilde{\mu}_{2,1}e^{i\theta_0}/\tilde{\mu}_{1,1}^{\frac{3}{2}} & \tilde{\mu}_{3,0}e^{i3\theta_0}/\tilde{\mu}_{1,1}^{\frac{3}{2}} & \\
& \tilde{\mu}_{0,4}e^{-i4\theta_0}/\tilde{\mu}_{1,1}^2 & 4\tilde{\mu}_{1,3}e^{-i2\theta_0}/\tilde{\mu}_{1,1}^2 & 6\tilde{\mu}_{2,2}/\tilde{\mu}_{1,1}^2 & 4\tilde{\mu}_{3,1}e^{i2\theta_0}/\tilde{\mu}_{1,1}^2 & \tilde{\mu}_{4,0}e^{i4\theta_0}/\tilde{\mu}_{1,1}^2 & \\
& & & \vdots & & & \\
\tilde{\mu}_{0,r}e^{-ir\theta_0}/\tilde{\mu}_{1,1}^{\frac{r}{2}} & \dots & \binom{r}{l} \tilde{\mu}_{l,r-l}e^{-i(r-2l)\theta_0}/\tilde{\mu}_{1,1}^{\frac{r}{2}} & \dots & \dots & \tilde{\mu}_{r,0}e^{ir\theta_0}/\tilde{\mu}_{1,1}^{\frac{r}{2}} &
\end{array}$$

where $\tilde{\mu}_{j,l}$ is the central complex moment and θ_0 satisfies (13).

5 Geometric Interpretation of the Moments

5.1 Shape Elongation

Intuitively, the elongation of a shape is described by the relationship between its length and its width. It is thus a property that is invariant under translation, scaling and rotation of the image. To characterize the shape elongation, we therefore consider the invariantization $\hat{m}'_2(\theta)$ of the second order moment $m_2(\theta)$ with respect to translation, scaling and rotation. By combining the results of Examples 1, 2 and 3, we have

$$\begin{aligned}
\hat{m}'_2(\theta) &= \frac{1}{4} \left(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} e^{i2\theta} + 2 + \frac{|\tilde{\mu}_{2,0}|}{\tilde{\mu}_{1,1}} e^{-i2\theta} \right), \\
&= \frac{1}{4} \left(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} (\cos(2\theta) + i\sin(2\theta)) + 2 + \frac{|\tilde{\mu}_{2,0}|}{\tilde{\mu}_{1,1}} (\cos(2\theta) - i\sin(2\theta)) \right), \\
&= \frac{1}{4} \left(2 \frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta) + 2 \right), \\
&= \frac{1}{2} \left(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta) + 1 \right), \tag{16}
\end{aligned}$$

where

$$\tilde{\mu}_{1,1} = \sum_{k=1}^N (z_k - z_0)(\bar{z}_k - \bar{z}_0) \rho(z_k) = \sum_{k=1}^N ((x_k - x_0)^2 + (y_k - y_0)^2) \rho(z_k) > 0.$$

Recall that when making the moments scaling invariant, we normalized $\hat{\mu}_{1,1}$ to 1 by dividing each $\tilde{\mu}_{j,l}$ by $\sqrt{\tilde{\mu}_{1,1}}$. Hence $\sqrt{\tilde{\mu}_{1,1}}$ represents the scale of the image.

Equation (16) indicates that the quantity $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}}$ prescribes the relationship between the maximum and the minimum values of the standard deviation of the random transform. It thus gives us a quantification of the elongation of the shape illustrated by the image⁴.

Lemma 8. *If not all $\rho(z_k)$ are zero, then $0 \leq \frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \leq 1$.*

Proof. By definition, $|\tilde{\mu}_{0,2}| \geq 0$ and $\tilde{\mu}_{1,1} > 0$, hence $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \geq 0$. If $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} > 1$, since $\hat{m}'_2(\theta) = \frac{1}{2}(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta) + 1)$, and $\hat{m}'_2(\theta) \geq 0$ by definition, if we choose $\theta = \frac{\pi}{2}$, then $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta) + 1 < 0$, which is a contradiction. Therefore $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \leq 1$. $\square$

The case $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 1$ corresponds to the most extreme elongation, namely the straight lines.

Lemma 9. *The pixel coordinates z_k lie on a single straight line if and only if $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 1$.*

Proof. $\Rightarrow$ Suppose we have a straight line. As the line is put vertically, the projection of the line is a dot. Hence the second moment of the radon transform is zero at that angle θ^* , i.e.

$$\hat{m}'_2(\theta^*) = \frac{1}{2}(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta^*) + 1) = 0. \quad (17)$$

Since $-1 \leq \cos(2\theta^*) \leq 1$ and from Lemma 8 we know that $0 \leq \frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \leq 1$ for any image, we can conclude that (17) is true only when $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 1$ and $\cos(2\theta^*) = -1$. Hence $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 1$ is true.

$\Leftarrow$ Now suppose $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 1$. Combine the results in Examples 1, 2 and 3, we conclude that

$$\hat{m}'_2(\theta) = \tilde{m}_2(\theta - \theta_0)/\tilde{\mu}_{1,1} = \frac{1}{2}(\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} \cos(2\theta) + 1), \quad \forall \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}],$$

where θ_0 satisfies (13). Since $\hat{m}'_2(\frac{\pi}{2}) = 0$, then there is $\tilde{\theta} = \frac{\pi}{2} - \theta_0$ such that at this angle the centralized second order moment $\tilde{m}_2(\tilde{\theta})$ of the image is zero.

Let the Radon transform $f_{\tilde{\theta}}(r)$ at angle $\tilde{\theta}$ be a discrete function. Without loss of generality, we assume that $\sum_{k=1}^N f_{\tilde{\theta}}(r_k(\theta)) = 1$ and $f_{\tilde{\theta}}(r_k(\theta)) \geq 0$. Since

$$\tilde{m}_2(\tilde{\theta}) = \sum_{k=1}^N (r_k(\tilde{\theta}) - r_0(\tilde{\theta}))^2 f_{\tilde{\theta}}(r_k(\theta)) = 0,$$

⁴A. R. Rostampour et al. [4] previously introduced the $\frac{n_{04}}{n_{02}^2}$ as a measure of elongation of the projections of an image. We observe that our measure of elongation has a lower order, and that it can be obtained without normalizing the angle of the image.

where $r_0(\tilde{\theta})$ is the projection of the centroid of the image, we observe that $f_{\tilde{\theta}}(r) = \delta(r - r_0(\tilde{\theta}))$. Then we conclude that the image lies on a line through $r_0(\tilde{\theta})$ with angle $\tilde{\theta} + \frac{\pi}{2}$ to the x-axis. $\square$

The other extreme case is when $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 0$. This corresponds to $\hat{m}'_2(\theta) = \text{constant}$, so the standard deviation of the projection is the same for all directions. There are many ways for this to happen. One interesting case is the discrete analogue of rotation symmetries.

Definition 2. An object is said to have $\tilde{N}$ -fold rotation symmetry ($\tilde{N}$ -FRS) if it is unchanged by a rotation around its centroid by $\frac{2k\pi}{\tilde{N}}$, for all $k = 1, \dots, \tilde{N}$.

Lemma 10. If the image I has an $\tilde{N}$ -FRS with $\tilde{N} > 2$, then $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 0$.

Proof. Suppose the data have $\tilde{N}$ -FRS. For a certain point with distance r_k , $k = 1, 2, \dots, M$, from the centroid, angle θ_k with x-axis and weight $\rho(z_k)$, there are $\tilde{N}_k - 1$ more points with the same distance from the centroid and the same weight $\rho(z_k)$ but having angle $\theta_k + \frac{2j\pi}{\tilde{N}_k}$, $j = 1, \dots, \tilde{N}_k - 1$ with x-axis respectively. Then the moment $\tilde{\mu}_{0,2}$ can be written as

$$\begin{aligned} \tilde{\mu}_{0,2} &= \sum_{k=1}^M \rho(z_k) \left(\sum_{j=0}^{\tilde{N}_k-1} \left(r_k \cos\left(\theta_k + \frac{2j\pi}{\tilde{N}_k}\right) - i r_k \sin\left(\theta_k + \frac{2j\pi}{\tilde{N}_k}\right) \right)^2 \right), \\ &= \sum_{k=1}^M \rho(z_k) \left(\sum_{j=0}^{\tilde{N}_k-1} r_k^2 \cos\left(2\theta_k + \frac{4j\pi}{\tilde{N}_k}\right) \right) - i \sum_{k=1}^M \rho(z_k) \left(\sum_{j=0}^{\tilde{N}_k-1} r_k^2 \sin\left(2\theta_k + \frac{4j\pi}{\tilde{N}_k}\right) \right), \\ &= \sum_{k=1}^M \rho(z_k) \left(\sum_{j=0}^{\tilde{N}_k-1} r_k^2 \cos(2\theta_k) \cos\left(\frac{4j\pi}{\tilde{N}_k}\right) - r_k^2 \sin(2\theta_k) \sin\left(\frac{4j\pi}{\tilde{N}_k}\right) \right) \\ &\quad - i \sum_{k=1}^M \rho(z_k) \left(\sum_{j=0}^{\tilde{N}_k-1} r_k^2 \sin(2\theta_k) \cos\left(\frac{4j\pi}{\tilde{N}_k}\right) + r_k^2 \cos(2\theta_k) \sin\left(\frac{4j\pi}{\tilde{N}_k}\right) \right), \\ &= \sum_{k=1}^M \rho(z_k) \left(r_k^2 \cos(2\theta_k) \left(\sum_{j=0}^{\tilde{N}_k-1} \cos\left(\frac{4j\pi}{\tilde{N}_k}\right) \right) - r_k^2 \sin(2\theta_k) \left(\sum_{j=0}^{\tilde{N}_k-1} \sin\left(\frac{4j\pi}{\tilde{N}_k}\right) \right) \right) \\ &\quad - i \sum_{k=1}^M \rho(z_k) \left(r_k^2 \sin(2\theta_k) \left(\sum_{j=0}^{\tilde{N}_k-1} \cos\left(\frac{4j\pi}{\tilde{N}_k}\right) \right) + r_k^2 \cos(2\theta_k) \left(\sum_{j=0}^{\tilde{N}_k-1} \sin\left(\frac{4j\pi}{\tilde{N}_k}\right) \right) \right). \end{aligned}$$

It can be shown that

$$\sum_{j=0}^{\tilde{N}_k-1} \cos\left(\frac{4j\pi}{\tilde{N}_k}\right) = 0, \quad \sum_{j=0}^{\tilde{N}_k-1} \sin\left(\frac{4j\pi}{\tilde{N}_k}\right) = 0, \quad \forall \tilde{N}_k > 2.$$

Then $\tilde{\mu}_{0,2} = 0$, hence $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = 0$. $\square$

One can give a statistical interpretation of our proposed shape elongation measure $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}}$. Indeed after renormalizing the total ink $\mu_{0,0} = \sum_{k=1}^N \rho(z_k)$ to one, one can view the pixel intensities as describing a discrete probability distribution. The standard deviation matrix of that distribution is then determined by the third row of the Pascal Triangle, as stated in the following Lemma:

Lemma 11. *Consider the discrete image I as a bivariate distribution with the joint probability mass function $P(X = x_k - x_0, Y = y_k - y_0) = \frac{\rho(z_k)}{\mu_{0,0}}$, $k = 1, 2, \dots, N$. The covariance matrix Σ of that distribution is given by*

$$\Sigma = \begin{pmatrix} \frac{\tilde{\mu}_{1,1} + \text{Re}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} & -\frac{\text{Im}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} \\ -\frac{\text{Im}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} & \frac{\tilde{\mu}_{1,1} - \text{Re}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} \end{pmatrix}.$$

Proof. Observe that

$$\begin{aligned} \frac{\tilde{\mu}_{1,1}}{\mu_{0,0}} &= \sum_{k=1}^N (z_k - z_0)(\bar{z}_k - \bar{z}_0) \frac{\rho(z_k)}{\mu_{0,0}} = \sum_{k=1}^N ((x_k - x_0)^2 + (y_k - y_0)^2) \frac{\rho(z_k)}{\mu_{0,0}}, \\ \frac{\tilde{\mu}_{0,2}}{\mu_{0,0}} &= \sum_{k=1}^N (\bar{z}_k - \bar{z}_0)^2 \frac{\rho(z_k)}{\mu_{0,0}} = \sum_{k=1}^N ((x_k - x_0)^2 - (y_k - y_0)^2) \frac{\rho(z_k)}{\mu_{0,0}} - 2i \sum_{k=1}^N (x_k - x_0)(y_k - y_0) \frac{\rho(z_k)}{\mu_{0,0}}. \end{aligned}$$

Write $\Sigma = \begin{pmatrix} \sigma_x^2 & \rho_{XY}\sigma_x\sigma_y \\ \rho_{XY}\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix}$. Since the random variables X, Y both have zero mean, we have

$$\sigma_x^2 = \sum_{k=1}^N (x_k - x_0)^2 \frac{\rho(z_k)}{\mu_{0,0}}, \quad \sigma_y^2 = \sum_{k=1}^N (y_k - y_0)^2 \frac{\rho(z_k)}{\mu_{0,0}}, \quad \rho_{XY}\sigma_x\sigma_y = \sum_{k=1}^N (x_k - x_0)(y_k - y_0) \frac{\rho(z_k)}{\mu_{0,0}}.$$

Then one can check that

$$\frac{\tilde{\mu}_{1,1}}{2\mu_{0,0}} + \frac{\text{Re}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} = \sigma_x^2, \quad -\frac{\text{Im}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} = \rho_{XY}\sigma_x\sigma_y, \quad \frac{\tilde{\mu}_{1,1}}{2\mu_{0,0}} - \frac{\text{Re}(\tilde{\mu}_{0,2})}{2\mu_{0,0}} = \sigma_y^2.$$

□

Recall that, in order to obtain the standard deviation $m_2(\theta)$ of the projection of a bivariate distribution onto the line with direction vector $(x, y) = r(\cos \theta, \sin \theta)$, one can simply project the standard deviation matrix Σ onto $(x, y) : (\cos \theta \quad \sin \theta) \Sigma \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = m_2(\theta)$.

It is easy to check that the relationship between the shape elongation $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}}$ and the eigenvalues $\lambda_{max}, \lambda_{min}$ of the standard deviation matrix Σ is $\frac{|\tilde{\mu}_{0,2}|}{\tilde{\mu}_{1,1}} = \left| \frac{\lambda_{max} - \lambda_{min}}{\lambda_{max} + \lambda_{min}} \right|$.

5.2 Rotational Symmetry

We have seen in the last section that an image I having an $\tilde{N}$ -FRS has $\tilde{\mu}_{0,2} = 0$. More generally, we have the following lemmas, which was used in [5] as the basis for a HAZMAT sign recognition method.

Lemma 12.⁵ *Let $\tilde{N}$ be a (finite) integer. If an image I has an $\tilde{N}$ -fold rotation symmetry and if $\frac{l-j}{\tilde{N}}$ is not an integer, then $\tilde{\mu}_{j,l} = 0$. Conversely, if $\tilde{\mu}_{j,l} = 0$ for all $\frac{l-j}{\tilde{N}}$ that are not an integer, then the image I has an $\tilde{N}$ -fold rotation symmetry.*

Proof. $\Rightarrow$ Let us rotate I clockwise around the origin by $\frac{2\pi}{\tilde{N}}$. Due to its symmetry, the rotated image I' must be the same as the original one. In particular, it must hold

$$\tilde{\mu}'_{j,l} = e^{2\pi i(l-j)/\tilde{N}} \tilde{\mu}_{j,l} = \tilde{\mu}_{j,l}.$$

Since $\frac{l-j}{\tilde{N}}$ is not an integer, this equation can be fulfilled only if $\tilde{\mu}_{j,l} = 0$.

$\Leftarrow$ Suppose $\tilde{\mu}_{j,0} = 0$ for any $\frac{j}{\tilde{N}}$ not an integer of some finite integer $\tilde{N}$. Let us rotate I clockwise around the origin by $\frac{2k\pi}{\tilde{N}}$ for each $k = 1, 2, \dots, \tilde{N} - 1$, then $\tilde{\mu}'_{j,0} = e^{-2\pi ijk/\tilde{N}} \tilde{\mu}_{j,0}$ for each k and all $j = 0, 1, \dots, \tilde{N} - 1$. For $\frac{jk}{\tilde{N}} \in \mathbb{Z}$, it is easy to check that $e^{-2\pi ijk/\tilde{N}} = 1$, hence $\tilde{\mu}'_{j,0} = \tilde{\mu}_{j,0}$. For $\frac{jk}{\tilde{N}} \notin \mathbb{Z}$, $\frac{j}{\tilde{N}}$ is not an integer either. Then $\tilde{\mu}_{j,0} = 0 = \tilde{\mu}'_{j,0}$. In this way we have $\tilde{\mu}'_{j,0} = \tilde{\mu}_{j,0}$ for all $j = 0, 1, \dots, \tilde{N} - 1$. Since in the proof of Lemma 1, we showed that $\{\tilde{\mu}_{j,0}\}_{j=0}^{\tilde{N}-1}$ uniquely determine I , then we can conclude that I and I' are the same for any rotation with angle $\frac{2k\pi}{\tilde{N}}$, $k = 1, 2, \dots, \tilde{N} - 1$. Therefore the image I has an $\tilde{N}$ -fold rotation symmetry. $\square$

Remark: As N increases, more and more columns of the Pascal Triangle $T^r(I)$ becomes zero, so that in the limit case, as $N \rightarrow \infty$, all entries $\mu_{j,l}$ with $j \neq l$ of $T^r(I)$, for any order r , vanish. This limit case corresponds to ∞ -fold rotation symmetry (e.g. circles), which however does not occur among discrete images.

5.3 Reflection Symmetry

Consider the reflection of an image about the line through the origin with direction vector $(\cos(\theta) \ \sin(\theta))^T$, $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2}]$. The point $z_k = x_k + iy_k$ is mapped to $\underline{z}_k = \bar{z}_k e^{i2\theta} = (x_k \cos(2\theta) + y_k \sin(2\theta)) + i(x_k \sin(2\theta) - y_k \cos(2\theta))$ under the reflection with its pixel inten-

⁵An analogue for the “if” part of this lemma for the case of a continuous image can be found in [6]

sity $\rho(z_k)$ staying the same. Then the new complex moment is

$$\begin{aligned}\underline{\mu}_{j,l} &= \sum_{k=1}^N \underline{z}_k^j \bar{\underline{z}}_k^l \rho(\underline{z}_k) = \sum_{k=1}^N (\bar{z}_k e^{i2\theta})^j (z_k e^{-i2\theta})^l \rho(z_k), \\ &= \sum_{k=1}^N e^{i2(j-l)\theta} \bar{z}_k^j z_k^l \rho(z_k) = e^{i2(j-l)\theta} \sum_{k=1}^N \bar{z}_k^j z_k^l \rho(z_k), \\ &= \mu_{l,j} e^{i2(j-l)\theta}, \quad \forall j, l \in \mathbb{Z}_{\geq 0}.\end{aligned}$$

Therefore the moment matrix for the new image $\underline{I}$ after reflection is

$$\tau_N(\underline{I}) = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{-i2\theta} & 0 & \cdots & 0 \\ 0 & 0 & e^{-i4\theta} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{-i2(N-1)\theta} \end{pmatrix} \tau_N(I)^T \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & e^{i2\theta} & 0 & \cdots & 0 \\ 0 & 0 & e^{i4\theta} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & e^{i2(N-1)\theta} \end{pmatrix}.$$

We can conclude from the above relation that if an image is symmetric with respect to the x-axis (i.e. $\theta = 0$), then we will have $\tau_N(I) = \tau_N(I)^T$, i.e. $\mu_{j,l} = \mu_{l,j}$ for all $j, l \in \mathbb{Z}_{\geq 0}$. Since $\mu_{j,l} = \bar{\mu}_{l,j}$ by definition, this means that all the $\mu_{j,l}$'s are real.

Similarly, if an image is symmetric with respect to the y-axis (i.e. $\theta = \frac{\pi}{2}$), we can conclude that $\mu_{j,l}$'s are real for j, l of the same parity and $\mu_{j,l}$'s are imaginary for j, l of opposite parity.

More generally, we have the following result:

Lemma 13. *A discrete image is symmetry with respect to reflections about a line through the origin with direction $(\cos \theta_0 \quad \sin \theta_0)^T$*

$$\iff \tan(l-j)\theta_0 = -\frac{\text{Im}(\mu_{j,l})}{\text{Re}(\mu_{j,l})}, \quad j, l = 0, 1, \dots$$

Proof. Notice that with reflection symmetry, we have

$$\begin{aligned}m_n(\theta_0 - \theta) &= \frac{1}{2^n} \sum_{k=1}^N (z_k e^{-i(\theta_0 - \theta)} + \bar{z}_k e^{i(\theta_0 - \theta)})^n \rho(z_k), \\ &= \frac{1}{2^n} \sum_{k=1}^N (\bar{z}_k e^{i2\theta_0} e^{-i(\theta_0 + \theta)} + z_k e^{-i2\theta_0} e^{i(\theta_0 + \theta)})^n \rho(z_k), \\ &= \frac{1}{2^n} \sum_{k=1}^N (z'_k e^{-i(\theta_0 + \theta)} + \bar{z}'_k e^{i(\theta_0 + \theta)})^n \rho(\bar{z}'_k e^{i2\theta_0}), \quad (\text{denote } z'_k = \bar{z}_k e^{i2\theta_0}) \\ &= \frac{1}{2^n} \sum_{k=1}^N (z'_k e^{-i(\theta_0 + \theta)} + \bar{z}'_k e^{i(\theta_0 + \theta)})^n \rho(z'_k), \\ &= m_n(\theta_0 + \theta).\end{aligned}$$

Conversely, if $m_n(\theta_0 - \theta) = m_n(\theta_0 + \theta)$ for an image $I = \{(z_k, \rho(z_k))\}_{k=1}^N$, then for its reflection image $I_r = \{(\bar{z}_k e^{i2\theta_0}, \rho(z_k))\}_{k=1}^N$, there is $m_n^r(\theta_0 + \theta) = m_n(\theta_0 - \theta) = m_n(\theta_0 + \theta)$ for any $\theta \in (-\pi, \pi]$. From Lemma 1 and Lemma 3 we can conclude that the image reconstructed from $\{m_n^r(\theta_j), j = 1, \dots, n+1\}_{n=0}^{N-1}$ and $\{m_n(\theta_j), j = 1, \dots, n+1\}_{n=0}^{N-1}$ will be the same, i.e. $I_r = I$. Hence the image has reflection symmetry.

By Equation (7), we know

$$m_n(\theta_0 + \theta) = \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \mu_{l,n-l} e^{i(n-2l)(\theta_0+\theta)}, \quad m_n(\theta_0 - \theta) = \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \mu_{l,n-l} e^{i(n-2l)(\theta_0-\theta)}.$$

Then $m_n(\theta_0 + \theta) = m_n(\theta_0 - \theta)$, i.e. $m_n(\theta_0 + \theta) - m_n(\theta_0 - \theta) = 0$ can be written as

$$\begin{aligned} & \frac{1}{2^n} \sum_{l=0}^n \binom{n}{l} \mu_{l,n-l} e^{i(n-2l)\theta_0} (e^{i(n-2l)\theta} - e^{-i(n-2l)\theta}) = 0, \\ \Leftrightarrow & \sum_{l=0}^n \binom{n}{l} \mu_{l,n-l} e^{i(n-2l)\theta_0} (2i \sin(n-2l)\theta) = 0, \\ \Leftrightarrow & \sum_{l=0}^n \binom{n}{l} \mu_{l,n-l} e^{i(n-2l)\theta_0} \sin(n-2l)\theta = 0, \\ \Leftrightarrow & \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{l} \sin(n-2l)\theta (\mu_{l,n-l} e^{i(n-2l)\theta_0} - \mu_{n-l,l} e^{i(2l-n)\theta_0}) = 0, \\ \Leftrightarrow & \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{l} \sin(n-2l)\theta 2i \operatorname{Im}(\mu_{l,n-l} e^{i(n-2l)\theta_0}) = 0, \\ \Leftrightarrow & \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{l} \sin(n-2l)\theta (\operatorname{Re}(\mu_{l,n-l}) \sin(n-2l)\theta_0 + \operatorname{Im}(\mu_{l,n-l}) \cos(n-2l)\theta_0) = 0. \end{aligned}$$

The above equation is true for any $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2}]$, so it is true if and only if

$$\operatorname{Re}(\mu_{l,n-l}) \sin(n-2l)\theta_0 + \operatorname{Im}(\mu_{l,n-l}) \cos(n-2l)\theta_0 = 0, \quad l = 0, 1, \dots, \lfloor \frac{n-1}{2} \rfloor,$$

i.e.

$$\tan(n-2l)\theta_0 = -\frac{\operatorname{Im}(\mu_{l,n-l})}{\operatorname{Re}(\mu_{l,n-l})}, \quad l = 0, 1, \dots, \lfloor \frac{n-1}{2} \rfloor. \quad (18)$$

Since $\mu_{n-l,l} = \bar{\mu}_{l,n-l}$, (18) can be written as

$$\tan(l-j)\theta_0 = -\frac{\operatorname{Im}(\mu_{j,l})}{\operatorname{Re}(\mu_{j,l})}, \quad \forall j = 0, 1, \dots, \quad l = 0, 1, \dots \quad (19)$$

□

In this way, for a given discrete image, we can check (19) to decide whether it is symmetric with respect to a certain line or not. If it is, then the line is at an angle θ given by

$$\theta = \begin{cases} \arctan\left(-\frac{\text{Im}(\mu_{j,j+1})}{\text{Re}(\mu_{j,j+1})}\right) & \text{if } \text{Re}(\mu_{j,j+1}) \neq 0, \forall j = 0, 1, \dots, \\ \frac{\pi}{2} & \text{otherwise.} \end{cases}$$

Acknowledgments

This research was supported in parts by NSF grant CCF-0728929. Address correspondence to Mireille Boutin (mboutin@purdue.edu).

References

- [1] Peyman Milanfar, George C. Verghese, W. Clem Karl and A. S. Willsky, “Reconstruction polygons from moments with connections to array processing,” *IEEE Transactions on Signal Processing* **43**, pp. 432–443, 1995.
- [2] Björn Gustafsson, Chiyu He, Peyman Milanfar and M. Putinar, “Reconstructing planar domains from their moments,” *Inverse Problems* **16**, pp. 1053–1070, 2000.
- [3] M. Fels and P. J. Olver, “Moving coframes: II. regularization and theoretical foundations,” *Acta Applicandae Mathematicae* **55**, pp. 127–208, 1999.
- [4] A. R. Rostampour and P. R. Madhupathy, “Shape recognition using simple measures of projections,” *Proceedings of 7th Intl. Phoenix Conference on Computers and Communications*, pp. 474–479, 1988.
- [5] Andrew W. Haddad, Shanshan Huang, Mireille Boutin and E. J. Delp, “Detection of symmetric shapes on a mobile device with applications to automatic sign interpretation,” *Proc. of SPIE-IS&T Electronic Imaging* **8304**, p. 83040G, 2012.
- [6] Jan Flusser, Barbara Zitova and T. Suk, *Moments and Moment Invariants in Pattern Recognition*, Wiley and Sons Ltd., 2009.
- [7] M. Fels and P. J. Olver, “Moving coframes: I. a practical algorithm,” *Acta Applicandae Mathematicae* **51**, pp. 161–213, 1998.
- [8] M.-K. Hu, “Visual pattern recognition by moment invariants,” *IRE Transactions on Information Theory* **8**, pp. 179–187, 1962.
- [9] P. J. Olver, *Classical Invariant Theory*, London Mathematical Society Student Texts, Cambridge University Press, 1999.

- [10] Peyman Milanfar, William C. Karl and A. S. Willsky, “A moment-based variational approach to tomographic reconstruction,” *IEEE Transactions on Image Processing* **5**, pp. 459–470, 1996.
- [11] M. P. Williams, “Solving polynomial equations using linear algebra,” *Johns Hopkins APL Technical Digest* **28**, pp. 354–363, 2010.